

THEORETICAL ASPECTS OF HIGGS PHYSICS[†]

BERND A. KNIEHL

*Max-Planck-Institut für Physik (Werner-Heisenberg-Institut),
Föhringer Ring 6, 80805 Munich, Germany**and**Institut für Theoretische Physik, Ludwig-Maximilians-Universität,
Theresienstraße 37, 80333 Munich, Germany*

We review the present status of Higgs physics within the standard model and its extensions. First, we briefly summarize the current experimental exclusion limits from the direct searches with LEP1 and the Tevatron, and assess the discovery potential of LEP2. Then, we report the mass bounds resulting from global fits to the latest electroweak precision data, and compile various theoretical principles which lead to restrictions—or even determinations—of the Higgs-boson masses. Perturbative upper bounds are discussed in some detail. Finally, we survey the recent progress in the computation of higher-order radiative corrections to the $b\bar{b}$ decay width of the standard-model Higgs boson.

1 Introduction

The $SU(2)_I \times U(1)_Y$ structure of the electroweak interactions has been consolidated by an enormous wealth of experimental data during the past three decades. The canonical way to generate masses for the fermions and intermediate bosons without violating this gauge symmetry in the Lagrangian is by the Higgs mechanism of spontaneous symmetry breaking. In the minimal standard model (SM), this is achieved by introducing one complex $SU(2)_I$ -doublet scalar field Φ with $Y = 1$. The three massless Goldstone bosons which emerge via the electroweak symmetry breaking are eaten up to become the longitudinal degrees of freedom of the $W^\pm$ and Z bosons, *i.e.*, to generate their masses, while one CP-even Higgs scalar boson H remains in the physical spectrum. The Higgs potential contains one mass and one self-coupling. Since the vacuum expectation value (VEV) is fixed by the relation $v = 2M_W/g$, where g is the $SU(2)_I$ gauge coupling, there remains one free parameter in the Higgs sector, namely, M_H .

A phenomenologically interesting extension of the SM Higgs sector that keeps the electroweak ρ parameter at unity in the Born approximation, is obtained by adding a second complex $SU(2)_I$ -doublet scalar field Φ_2 with $Y = -1$. This leads to the two-Higgs-doublet model (2HDM). After the three Goldstone bosons have been eliminated, there remain five physical Higgs scalars: the neutral CP-even h^0 and H^0 bosons, the neutral CP-odd A^0 boson, and the charged $H^\pm$ pair. The

most general CP-conserving Higgs potential for the 2HDM contains three masses and four self-couplings. Subtracting the M_W constraint on the two VEV's v_1 and v_2 , one is left with six free parameters, which are usually taken to be m_{h^0} , m_{H^0} , m_{A^0} , $m_{H^\pm}$, $\tan\beta = v_2/v_1$, and the weak mixing angle α which relates the weak and mass eigenstates of h^0 and H^0 . According to the Glashow-Weinberg theorem,¹ flavour-changing neutral currents may be avoided if all fermions with the same electric charge couple to the same Higgs doublet. In the 2HDM of type II, the up-type/down-type fermions couple to the Higgs doublet with $Y = \pm 1$.

The Higgs sector of the minimal supersymmetric extension of the SM (MSSM) consists of a 2HDM of type II. Since the Higgs self-couplings are then determined by the gauge couplings, there are only two free parameters in the MSSM Higgs sector, which are conveniently chosen to be m_{A^0} and $\tan\beta$. However, a large number of new masses, couplings, and mixing angles connected with the supersymmetric partners enter via loop effects. In the supergravity-inspired MSSM, all these degrees of freedom may be related to just five parameters at the GUT scale: the Higgs mixing μ , the sfermion mass m_0 , the gaugino mass $m_{1/2}$, the trilinear sfermion-Higgs coupling A , and the bilinear Higgs coupling B . After the renormalization-group (RG) evolution down to the electroweak scale, the heavy sfermions are approximately degenerate, with mass M_S . The following three scenarios are often considered in the literature: (i) no mixing: $A = 0$, $|\mu| \ll M_S$; (ii) typical mix-

[†] To appear in the *Proceedings of the 28th International Conference on High Energy Physics*, Warsaw, Poland, 25–31 July 1996, edited by A. Wróblewski.

Table 1: 95% CL M_H lower bounds (in GeV) from LEP1.

Collab.	Years	$10^6 q\bar{q}$	$M_H^{\min}$
ALEPH	89–95	4.5	63.9
DELPHI	91, 92, 94, 95	3.1	58.3
L3	91–94	3.1	60.2
OPAL	89–95	4.4	59.6

ing: $A = -\mu = M_S$; and (iii) maximal mixing: $A = \sqrt{6}M_S$, $|\mu| \ll M_S$.

2 Direct Higgs searches

2.1 LEP1 and Tevatron

The SM Higgs boson was searched at LEP1 via Bjorken's process $e^+e^- \rightarrow Hf\bar{f}$ by looking for a pair of acoplanar jets together with missing energy in the $H\nu\bar{\nu}$ channel, or for a hadronic event with an energetic lepton pair in the HL^+l^- ($l = e, \mu$) channel. The extraction by ALEPH of the 95% CL lower bound on M_H is illustrated in Fig. 1. The present LEP1 bounds ^{2,3} are summarized in Table 1.




Figure 1: M_H dependence of the number of Higgs signal events expected by ALEPH.

The h^0 and A^0 bosons of the 2HDM or MSSM could have been produced at LEP1 via Higgsstrahlung $e^+e^- \rightarrow h^0 f\bar{f}$, where the h^0 boson is radiated off the resonant Z boson, associated

Table 2: 95% CL lower bounds on m_{h^0} , m_{A^0} , and $m_{H^\pm}$ (in GeV) from LEP1.

Collab.	$m_{h^0}^{\min}$	$m_{A^0}^{\min}$	$m_{H^\pm}^{\min}$
ALEPH	45.5	45	41.7
DELPHI	45.4	45.2	—
L3	42	23	42.8
OPAL	44.3	23.5	44.1

production $e^+e^- \rightarrow h^0 A^0$, and the Yukawa process $e^+e^- \rightarrow h^0/A^0 f\bar{f}$ ($f = b, \tau$), where the h^0 and A^0 bosons couple to the fermion line. The $h^0 Z^*$ and $h^0 A^0$ channels are complementary, in the sense that their cross sections are proportional to $\sin^2(\beta - \alpha)$ and $\cos^2(\beta - \alpha)$, respectively. In the framework of the MSSM with $M_S = 1$ TeV and no squark mixing, DELPHI has excluded the shaded regions in the $(m_{h^0}, \tan\beta)$ plane of Fig. 2. The current 95% CL lower bounds on m_{h^0} and m_{A^0} in the MSSM obtained by the four LEP1 experiments ^{3,4} are listed in Table 2.




Figure 2: Region in the $(m_{h^0}, \tan\beta)$ plane excluded at 95% CL by DELPHI. The dark areas are forbidden by theory.

In the more general context of the 2HDM, the h^0 (A^0) boson could have been missed in the $h^0 Z^*$ and $h^0 A^0$ channels, if $\sin^2(\beta - \alpha) \approx 0$ (1) and (or) $m_{h^0} + m_{A^0} > M_Z$. Then, the Yukawa process would be dominant.⁵ The MSSM scenario with $m_{A^0} \ll M_Z$ and $\tan\beta \gg 1$ was considered particularly interesting⁶ since it could par-

tially explain the R_b anomaly, which was then still existing. ALEPH⁷ searched for the h^0 and A^0 bosons of the 2HDM using the Yukawa process with $b\bar{b}b\bar{b}$, $\tau^+\tau^-b\bar{b}$, $\tau^+\tau^-\tau^+\tau^-$, and $\tau^+\tau^-X$ final states, where X is a system with low charged multiplicity. The $b\bar{b}b\bar{b}$ channel turned out to be less efficient in ruling out the small- m_{A^0} large- $\tan\beta$ scenario than anticipated.⁶ It is amusing to observe that, in the 2HDM with $\alpha \approx \beta$, the most stringent m_{h^0} lower bound presently comes from HERA.⁸ The future E821 experiment at BNL is expected to decrease the experimental error on $(g-2)_\mu$ by a factor of twenty or more,⁹ and will thus lead to interesting m_{h^0} and m_{A^0} lower bounds via loop effects.¹⁰

Searching for the $H^\pm$ bosons of the 2HDM via pair production $e^+e^- \rightarrow H^+H^-$, the LEP1 experiments¹¹ could almost exclude the entire $m_{H^\pm}$ range allowed by kinematics (see Table 2). CDF¹² has looked for $H^\pm$ bosons in the decay products of $t\bar{t}$ pairs, and has greatly increased the LEP1 lower bound on $m_{H^\pm}$ for $\tan\beta \gtrsim 60$ (see Fig. 3). If $\tan\beta \gg 1$, then the cascade $t \rightarrow H^+b \rightarrow \tau^+\nu_\tau b$ is more likely to happen than $t \rightarrow W^+b \rightarrow \tau^+\nu_\tau b$ and leads to a larger missing transverse energy.¹²




Figure 3: Region in the $(\tan\beta, m_{H^\pm})$ plane excluded at 95% CL by CDF.

2.2 LEP2

The theoretical and experimental aspects of Higgs phenomenology at LEP2 have recently been summarized in a comprehensive report.¹³ Higgs-strahlung is the dominant production mechanism of the SM Higgs boson at LEP2. In Fig. 4, its cross section is shown as a function of M_H for three values of CM energy.¹⁴ Electroweak radiative corrections¹⁵ are included here.




Figure 4: Cross section of Higgs-strahlung at LEP2 as a function of M_H .

The $H\nu_e\bar{\nu}_e$ and He^+e^- final states are also produced through W^+W^- and ZZ fusion, respectively. In the threshold region, the signal cross section is appreciably increased by coherently including these processes.¹⁶ This is illustrated for $H\nu_e\bar{\nu}_e$ production in Fig. 5.




Figure 5: $H\nu_e\bar{\nu}_e$ production via Higgs-strahlung, W^+W^- fusion, and their interference in the threshold region at LEP2.

The minimum luminosity $L_{\min}$ needed per experiment for a combined 5σ discovery or a 95% CL exclusion of the SM Higgs boson under realistic LEP2 conditions¹³ is shown as a function of M_H in Fig. 6. We see that, at $\sqrt{s} = 192$ GeV, $L_{\min} = 150 \text{ pb}^{-1}$ is sufficient to discover the SM Higgs boson with $M_H \lesssim 95$ GeV. On the other hand, a 95 GeV Higgs boson can be excluded at

95% CL with $L_{\min}$ as low as 33 pb^{-1} , while, with $L_{\min} = 150 \text{ pb}^{-1}$, the M_H range way up to 98 GeV may be excluded.




Figure 6: LEP2 exclusion and discovery limits for the SM Higgs boson.

The 5σ discovery and 95% CL exclusion limits in the $(m_{h^0}, \tan\beta)$ plane of the MSSM that may be reached at LEP2 with a luminosity of 150 pb^{-1} per experiment¹³ may be seen from Fig. 7. Here, $M_t = 175 \text{ GeV}$, $M_S = 1 \text{ TeV}$, and typical squark mixing as described in the Introduction are assumed. Comparing Figs. 2 and 7, we observe that LEP2 will be able to close the low- $\tan\beta$ region which was not covered by LEP1.




Figure 7: LEP2 exclusion and discovery limits in the $(m_{h^0}, \tan\beta)$ plane.

3 Higgs mass bounds

3.1 Global fits

Experimental precision tests of the standard electroweak theory are sensitive to the Higgs boson via quantum corrections. A global fit¹⁷ to the e^+e^- data on the Z-boson observables from ALEPH, DELPHI, L3, OPAL, and SLD, the $p\bar{p}$ data on M_W from CDF, D0, and UA2, the νN data on $\sin^2\theta_w$ from CCFR, CDHS, and CHARM, and the Tevatron data on M_t from CDF and D0 which were presented during this conference yields $M_H = 149^{+148}_{-82} \text{ GeV}$ with $\chi^2_{\min}/\text{d.o.f.} = 19/14 \approx 1.36$. The resulting M_H distribution of $\Delta\chi^2 = \chi^2 - \chi^2_{\min}$ is shown in Fig. 8, where the shaded band represents the estimated¹⁷ theoretical error due to missing higher-order corrections. From Fig. 8, one may read off an 95% CL upper bound on M_H at 550 GeV.¹⁷ The electroweak precision data have reached a quality which even allows for global fits within the MSSM.¹⁸




Figure 8: M_H dependence of $\Delta\chi^2 = \chi^2 - \chi^2_{\min}$ resulting from a global fit to the latest experimental data.

3.2 Theoretical principles

There have been various attempts to bound or even determine M_H from first principles.

Naturalness

By requiring that the Higgs-boson mass counterterm, *i.e.*, the difference between the bare and renormalized masses, be devoid of quadratic ultraviolet divergences, Veltman derived the mass relation¹⁹

$$\sum_f N_f M_f^2 = \frac{3}{2} M_W^2 + \frac{3}{4} M_Z^2 + \frac{3}{4} M_H^2, \quad (1)$$

where $N_f = 1$ (3) for leptons (quarks). Naïvely inserting the known pole masses, this leads to $M_H \approx 320$ GeV. Since this prediction is based on fine-tuning, it is nowadays considered unnatural.

Noncommutative geometry

It is possible²⁰ to construct the SM on the basis of the graded Lie algebra $SU(2|1)$, which contains the $SU(2)_I \times U(1)_Y$ Lie algebra in its even part. The essential ingredient is an algebraic superconnection which incorporates both the gauge and Higgs fields and whose curvature automatically generates the spontaneously broken realization of the SM. This leads to a geometrical interpretation of the Higgs mechanism. As a by-product, one obtains²⁰ the mass relation $M_H = \sqrt{2} M_W$. Taking this as an initial condition at the GUT scale and performing a one-loop RG analysis, one finds²¹ the physical Higgs-boson mass to be $M_H = 164$ GeV if $M_t = 175$ GeV.

Triviality and vacuum stability

Triviality²² and vacuum stability²³ are features connected with the RG-improved effective potential $V_{\text{eff}} = -m^2(\mu)|\Phi|^2 + \lambda(\mu)|\Phi|^4/2$. The dependence of the quartic self-coupling $\lambda(\mu)$ on the renormalization scale μ is determined by the RG equations. Roughly speaking, the triviality upper bound (vacuum-stability lower bound) on M_H follows from the requirement that $\lambda(\mu)$ stay finite (positive) for all $\mu < \Lambda$, where Λ is the cutoff beyond which new physics operates. Assuming the SM to be valid up to the GUT scale $\Lambda = 10^{16}$ GeV, one thus obtains¹³ $130 \text{ GeV} \lesssim M_H \lesssim 180 \text{ GeV}$. In turn, should the SM Higgs boson be discovered at LEP2, new physics is expected to appear below $\Lambda = 10 \text{ TeV}$.

Metastability

Depending on M_H and M_t , V_{eff} at finite (or zero) temperature may have a deep stable minimum at $\langle |\Phi_{\text{min}}| \rangle \gg G_F^{-1/2}$, *i.e.*, the physical electroweak minimum may just be metastable. Then, an absolute lower bound on M_H follows from the condition that the probability, normalized w.r.t. the expansion rate of the Universe, for the decay of the metastable vacuum by thermal fluctuations (or quantum tunneling) be negligibly small.²⁴ The M_H bounds thus obtained are somewhat below the usual stability bounds.²³

Multiple point criticality principle

It has been argued²⁵ that a mild form of locality breaking in quantum gravity due to baby universes, which are expected to render coupling constants dynamical, leads to the realization of the so-called multiple-point criticality principle in Nature. According to this principle, Nature should choose coupling constants such that the vacuum can exist in degenerate phases, *i.e.*, $V_{\text{eff}}(\Phi_{\text{min},1}) = V_{\text{eff}}(\Phi_{\text{min},2})$. In order that the dynamical mechanism be relevant, these authors²⁵ also require a strong first-order phase transition between the two vacua, implemented by $\langle |\Phi_{\text{min},2}| \rangle \approx M_{\text{Planck}} = 2 \times 10^{19} \text{ GeV}$. Via the usual RG analysis, these two assumptions then lead to a simultaneous prediction of M_t and M_H , namely,²⁵ $M_t = (173 \pm 4) \text{ GeV}$ and $M_H = (135 \pm 9) \text{ GeV}$.

Perturbation-theory breakdown

An attractive way of constraining M_H from above is to require that the Higgs sector be weakly interacting, so that perturbation theory is meaningful.²⁶ The resulting bounds depend somewhat on the considered process and the precise definition of perturbation-theory breakdown, but they are independent of assumptions concerning the scale Λ of new physics. At one and two loops, the leading high- M_H corrections to physical observables related to Higgs-boson production or decay are of $\mathcal{O}(G_F M_H^2)$ and $\mathcal{O}(G_F^2 M_H^4)$, respectively. For M_H increasing, the $\mathcal{O}(G_F^2 M_H^4)$ corrections eventually exceed the $\mathcal{O}(G_F M_H^2)$ ones in size, so that the perturbative expansions in $G_F M_H^2$ cease to usefully converge. The values M_H^{max} where this happens may be used to define a perturbative upper bound on M_H .

As first examples, the Higgs decays to pairs of fermions²⁷ and intermediate bosons²⁸ have recently been studied through $\mathcal{O}(G_F^2 M_H^4)$. This task may be greatly simplified in the limit of interest, $M_H \gg 2M_Z$, through the use of the Goldstone-boson equivalence theorem.²⁹ This theorem states that the leading high- M_H electroweak contribution to a Feynman diagram may be calculated by replacing the intermediate bosons $W^\pm$ and Z with the respective would-be Goldstone bosons $w^\pm$ and z of the symmetry-breaking sector. In this limit, the gauge and Yukawa couplings may be neglected against the Higgs self-coupling. By the same token, the Goldstone bosons may be taken to be massless, and the fermion loops may be omitted. The resulting correction factor K_f for $\Gamma(H \rightarrow f\bar{f})$ is independent of the fermion flavour f . Similarly, $\Gamma(H \rightarrow W^+W^-)$ and $\Gamma(H \rightarrow ZZ)$ receive the same correction factor K_V . In the on-mass-shell renormalization scheme, the results read^{27,28}

$$\begin{aligned}
K_f &= 1 + \hat{\lambda} \left(13 - 2\pi\sqrt{3} \right) + \hat{\lambda}^2 \left[12 - 169\pi\sqrt{3} \right. \\
&\quad \left. + 170\zeta(2) - 252\zeta(3) + 12 \left(13\pi + 19\sqrt{3} \right) \right. \\
&\quad \left. \times \text{Cl}_2 \left(\frac{\pi}{3} \right) \right] \\
&\approx 1 + 11.1\% \left(\frac{M_H}{\text{TeV}} \right)^2 - 8.9\% \left(\frac{M_H}{\text{TeV}} \right)^4, \\
K_V &= 1 + \hat{\lambda} \left(19 - 6\pi\sqrt{3} - 10\zeta(2) \right) + 62.0 \hat{\lambda}^2 \\
&\approx 1 + 14.6\% \left(\frac{M_H}{\text{TeV}} \right)^2 + 16.9\% \left(\frac{M_H}{\text{TeV}} \right)^4, \quad (2)
\end{aligned}$$

where $\hat{\lambda} = (G_F M_H^2 / 16\pi^2 \sqrt{2})$, ζ is Riemann's zeta function, and Cl_2 is Clausen's integral. The $\mathcal{O}(G_F^2 M_H^2)$ terms in Eq. (2) have been known for a long time.³⁰ K_f and K_V are displayed as functions of M_H in Fig. 9, from where we may read off the M_H^{max} values 1114 GeV and 930 GeV, respectively. The nonperturbative value of K_V at $M_H = 727$ GeV may be extracted from a recent lattice simulation of elastic $\pi\pi$ scattering in the framework of the four-dimensional O(4)-symmetric nonlinear σ model in the broken phase, where the σ resonance was observed.³¹

The study of the μ dependence of Higgs-boson observables in the $\overline{\text{MS}}$ renormalization scheme provides another aspect of perturbation-theory breakdown in the Higgs sector. If perturbation theory is valid, the μ dependence should be reduced as




Figure 9: K_V and K_f to $\mathcal{O}(G_F^2 M_H^2)$ and $\mathcal{O}(G_F^2 M_H^4)$ as functions of M_H . The crosses indicate the tree-level, one-loop, two-loop, and nonperturbative values of K_V at $M_H = 727$ GeV.

higher-order corrections are included. While this empirical rule is satisfied for $\Gamma(H \rightarrow W^+W^-)$ at $M_H = 400$ GeV, it is clearly violated at $M_H = 700$ GeV,³² as may be seen from Fig. 10.




Figure 10: μ dependence of K_V in the $\overline{\text{MS}}$ scheme at $M_H = 400$ and 700 GeV.

4 Radiative corrections to $\Gamma(H \rightarrow b\bar{b})$

The SM Higgs boson with intermediate mass $M_H \lesssim 2M_W$ decays dominantly to $b\bar{b}$ pairs. The radiative corrections to the partial width of this decay involve two very different scales, namely, M_H and M_t . It is therefore convenient to treat this process in the framework of a $n_f = 5$ effective Yukawa Lagrangian, *i.e.*, to integrate out the top quark. This leads to a RG-improved formulation,

which provides a natural separation of the $n_f = 5$ QCD corrections at scale $\mu = M_H$ and the top-quark-induced $n_f = 6$ corrections at scale $\mu = M_t$. One thus obtains³³ the following structure:

$$\Gamma_{b\bar{b}} = \Gamma_{b\bar{b}}^{\text{Born}} \left[\left(1 + \Delta_b^{\text{QED}}\right) \left(1 + \Delta_b^{\text{weak}}|_{x_t=0}\right) \times \left(1 + \Delta_b^{\text{QCD}}\right) \left(1 + \Delta_b^t\right) + \Xi_b^{\text{QCD}} \Xi_b^t \right]. \quad (3)$$

If the Born formula

$$\Gamma_{b\bar{b}}^{\text{Born}} = \frac{3G_F M_H m_b^2}{4\pi\sqrt{2}} \left(1 - \frac{4m_b^2}{M_H^2}\right)^{3/2} \quad (4)$$

is written with the QED and QCD $\overline{\text{MS}}$ mass evaluated with $n_f = 5$ quark flavours at scale $\mu = M_H$, $m_b^{(5)}(M_H)$, then Δ_b^{QED} and Δ_b^{QCD} are finite for $m_b = 0$ and read³⁴

$$\begin{aligned} \Delta_b^{\text{QED}} &= \frac{17}{36} \frac{\alpha}{\pi}, \\ \Delta_b^{\text{QCD}} &= \frac{17}{3} a_5 + a_5^2 \left(\frac{8851}{144} - \frac{47}{6} \zeta(2) - \frac{97}{6} \zeta(3) \right) \\ &\quad + a_5^3 \left(\frac{34873057}{46656} - \frac{10225}{54} \zeta(2) \right. \\ &\quad \left. - \frac{80095}{216} \zeta(3) - \frac{25}{6} \zeta(4) + \frac{1945}{36} \zeta(5) \right) \\ &= 5.66667 a_5 + 29.14671 a_5^2 + 41.75760 a_5^3, \end{aligned} \quad (5)$$

where $a_5 = \alpha_s^{(5)}(M_H)/\pi$. By means of scale optimization according to the principles of fastest apparent convergence (FAC) or minimal sensitivity (PMS), the coefficient of a_5^4 may be estimated³⁵ to be -981 . In the limit $M_H \ll 2M_W$, the weak correction, with the $\mathcal{O}(G_F M_t^2)$ term stripped off, reads³⁶

$$\Delta_b^{\text{weak}}|_{x_t=0} = \frac{G_F M_Z^2}{8\pi^2\sqrt{2}} \left(\frac{1}{6} - \frac{7}{3} c_w^2 - \frac{16}{3} c_w^4 + 3 \frac{c_w^2}{s_w^2} \ln c_w^2 \right), \quad (6)$$

where $c_w^2 = 1 - s_w^2 = M_W^2/M_Z^2$. The leading contributions due to the $b\bar{b}$ and $b\bar{b}g$ cuts of the double-triangle diagrams where the top quark circulates in one of the triangles are contained in^{33,37}

$$\Xi_b^{\text{QCD}} = a_5 \left(-\frac{76}{3} + 8\zeta(2) - \frac{4}{3} \ln^2 \frac{m_b^2}{M_H^2} \right). \quad (7)$$

The would-be mass singularity proportional to $\ln^2(m_b^2/M_H^2)$ cancels if the $b\bar{b}(g)$ and gg decay

channels are combined.³⁷ The leading top-quark-induced corrections are concentrated in³³

$$\begin{aligned} \Delta_b^t &= a_6^2 \left(\frac{5}{9} + \frac{2}{3} L \right) + x_t \left[1 + a_6 \left(-\frac{4}{3} - 4\zeta(2) \right) \right. \\ &\quad \left. + a_6^2 \left(-59.16260 + \frac{2}{3} L \right) \right], \\ \Xi_b^t &= a_6 \left(-\frac{1}{12} - \frac{1}{12} x_t \right), \end{aligned} \quad (8)$$

where $\mu_t = m_t^{(6)}(\mu_t)$, $a_6 = \alpha_s^{(6)}(\mu_t)/\pi$, $L = \ln(\mu_t^2/M_H^2)$, and $x_t = (G_F \mu_t^2/8\pi^2\sqrt{2})$. The L -dependent terms may be resummed by exploiting the RG-invariance of the energy-momentum tensor.³³

The QCD correction³⁸ to $\Gamma(H \rightarrow gg)$ includes a contribution due to the $b\bar{b}g$ final state, which may also be interpreted³⁹ as a $\mathcal{O}(\alpha_s^3 M_H^2/m_b^2)$ correction to $\Gamma(H \rightarrow b\bar{b})$.

Acknowledgments

The author is indebted to Ralf Hempfling for beneficial discussions.

References

1. S.L. Glashow and S. Weinberg, *Phys. Rev. D* **15**, 1958 (1977).
2. D. Buskulic *et al.* (ALEPH Collaboration), Pa13-026, *Phys. Lett. B* **384**, 427 (1996); M. Acciarri *et al.* (L3 Collaboration), Pa11-016, *Phys. Lett. B* **385**, 454 (1996).
3. C. Martínez Rivero *et al.* (DELPHI Collaboration), Pa07-095; P. Igo-Kemenes *et al.* (OPAL Collaboration), Pa13-004.
4. P. Janot, *Nucl. Phys. B (Proc. Suppl.)* **38**, 264 (1995); O. Adriani *et al.* (L3 Collaboration), *Z. Phys. C* **57**, 355 (1993).
5. J. Kalinowski and M. Krawczyk, *Phys. Lett. B* **361**, 66 (1995); *Acta Phys. Pol. B* **27**, 961 (1996).
6. J.D. Wells, C. Kolda, and G.L. Kane, *Phys. Lett. B* **338**, 219 (1994); *Phys. Rev. Lett.* **76**, 869 (1996).
7. ALEPH Collaboration, Pa13-027.
8. A.C. Bawa and M. Krawczyk, *Phys. Lett. B* **357**, 637 (1995).
9. L. Roberts, in these proceedings.

10. M. Krawczyk and J. Żochowski, Report Nos. IFT 15/96 and hep-ph/9608321 (August 1996).
11. D. Decamp *et al.* (ALEPH Collaboration), *Phys. Rep.* **216**, 253 (1992); B. Lindemann and A. Sopczak, L3 Note Nos. 1966 and hep-ph/9607327 (June 1996); G. Alexander *et al.* (OPAL Collaboration), *Phys. Lett. B* **370**, 174 (1996).
12. F. Abe *et al.* (CDF Collaboration), *Phys. Rev. D* **54**, 735 (1996); Pa13-025.
13. M. Carena, P.M. Zerwas (conveners) *et al.*, in *Physics at LEP2*, edited by G. Altarelli, T. Sjöstrand, and F. Zwirner, CERN Yellow Report No. 96-01, Vol. 1, p. 351.
14. E. Gross, B.A. Kniehl, and G. Wolf, *Z. Phys. C* **63**, 417 (1994); **66**, 321(E) (1995).
15. J. Fleischer and F. Jegerlehner, *Nucl. Phys. B* **216**, 469 (1983); F.A. Berends and R. Kleiss, *Nucl. Phys. B* **260**, 32 (1985); B.A. Kniehl, *Z. Phys. C* **55**, 605 (1992); A. Denner, J. Küblbeck, R. Mertig, and M. Böhm, *Z. Phys. C* **56**, 261 (1992).
16. W. Kilian, M. Krämer, and P.M. Zerwas, *Phys. Lett. B* **373**, 135 (1996).
17. J. Alcaraz *et al.* (LEP Electroweak Working Group and SLD Heavy Flavor Group), CERN Report No. LEPEWWG/96-02 (30 July 1996).
18. W. de Boer, A. Dabelstein, W. Hollik, W. Möhle, and U. Schwickerath, Report Nos. Pa11-001, IEKP-KA/96-08, KA-TP-18-96, and hep-ph/9609209 (August 1996).
19. M. Veltman, *Acta Phys. Pol. B* **12**, 437 (1981).
20. R. Häußling, N.A. Papadopoulos, and F. Scheck, *Phys. Lett. B* **260**, 125 (1991); R. Coquereaux, R. Häußling, N.A. Papadopoulos, and F. Scheck, *Int. J. Mod. Phys. A* **7**, 2809 (1992).
21. Y. Okumura, Report Nos. CHUBU9607 and hep-ph/9608208 (October 1996).
22. N. Cabibbo, L. Maiani, G. Parisi, and R. Petronzio, *Nucl. Phys. B* **158**, 295 (1979); M. Lindner, *Z. Phys. C* **31**, 295 (1986).
23. G. Altarelli and G. Isidori, *Phys. Lett. B* **337**, 141 (1994); J.A. Casas, J.R. Espinosa, and M. Quirós, *Phys. Lett. B* **342**, 171 (1995), *Phys. Lett. B* **382**, 374 (1996).
24. J.R. Espinosa and M. Quirós, *Phys. Lett. B* **353**, 257 (1995).
25. C.D. Froggatt and H.B. Nielsen, Report Nos. GUTPA-96-7-2, Pa08-012, and hep-ph/9607302 (July 1996); see also R. Rodenberger, Pa12-001 (July 1996).
26. M. Veltman, *Phys. Lett. B* **70**, 253 (1977).
27. L. Durand, B.A. Kniehl, and K. Riesselmann, *Phys. Rev. Lett.* **72**, 2534 (1994); **74**, 1699(E) (1995); A. Ghinculov, *Phys. Lett. B* **337**, 137 (1994); **346**, 426(E) (1995); V. Borodulin and G. Jikia, Report No. Freiburg-THEP 96/19 and hep-ph/9609447 (September 1996).
28. A. Ghinculov, *Nucl. Phys. B* **455**, 21 (1995); A. Frink, B.A. Kniehl, D. Kreimer, and K. Riesselmann, Pa07-081, *Phys. Rev. D* **54**, 4548 (1996).
29. J.M. Cornwall, D.N. Levin, and G. Tiktopoulos, *Phys. Rev. D* **10**, 1145 (1974); **11**, 972(E) (1975); C.E. Vayonakis, *Lett. Nuovo Cimento* **17**, 383 (1976).
30. M. Veltman, *Acta Phys. Pol. B* **8**, 475 (1977); W.J. Marciano and S.S.D. Willenbrock, *Phys. Rev. D* **37**, 2509 (1988).
31. M. Göckeler, H.A. Kastrup, J. Westphalen, and F. Zimmermann, *Nucl. Phys. B* **425**, 413 (1994).
32. U. Nierste and K. Riesselmann, *Phys. Rev. D* **53**, 6638 (1996).
33. K.G. Chetyrkin, B.A. Kniehl, and M. Steinhauser, Report Nos. MPI/PhT/96-65 and hep-ph/9610456 (July 1996).
34. E. Braaten and J.P. Leveille, *Phys. Rev. D* **22**, 715 (1980); S.G. Gorishny, A.L. Kataev, S.A. Larin, and L.R. Surguladze, *Mod. Phys. Lett. A* **5**, 2703 (1990); K.G. Chetyrkin, Report Nos. MPI/PhT/96-61 and hep-ph/9608318 (August 1996), *Phys. Lett. B* (in press).
35. K.G. Chetyrkin, B.A. Kniehl, and A. Sirlin, Report No. MPI/PhT/96-62.
36. B.A. Kniehl, *Nucl. Phys. B* **376**, 3 (1992).
37. S.A. Larin, T. van Ritbergen, and J.A.M. Vermaseren, *Phys. Lett. B* **362**, 134 (1995); K.G. Chetyrkin and A. Kwiatkowski, *Nucl. Phys. B* **461**, 3 (1996).
38. T. Inami, T. Kubota, and Y. Okada, *Z. Phys. C* **18**, 69 (1983); A. Djouadi, M. Spira, and P.M. Zerwas, *Phys. Lett. B* **264**, 440 (1991).
39. A. Djouadi, M. Spira, and P.M. Zerwas, *Z. Phys. C* **70**, 427 (1996).